

High heart rate reactors display greater decreases in tear SIgA concentration following a novel acute stressor

Hanstock, Helen; Edwards, Jason; Roberts, Ross; Walsh, Neil

Biological Psychology

DOI:

[10.1016/j.biopsycho.2018.02.002](https://doi.org/10.1016/j.biopsycho.2018.02.002)

Published: 01/03/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Hanstock, H., Edwards, J., Roberts, R., & Walsh, N. (2018). High heart rate reactors display greater decreases in tear SIgA concentration following a novel acute stressor. *Biological Psychology*, 133(March), 85-88. <https://doi.org/10.1016/j.biopsycho.2018.02.002>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**High heart rate reactors display greater decreases in tear SIgA concentration following
a novel acute stressor**

Running header: Stress reactivity and tear SIgA response to stress.

Helen G. Hanstock^{a,b}, Jason P. Edwards^a, Ross Roberts^c and Neil P. Walsh^a

^a Extremes Research Group, College of Health and Behavioural Sciences, Bangor University,
Bangor, Gwynedd, UK.

^b Swedish Winter Sports Research Centre, Department of Health Sciences, Mid Sweden
University, Östersund, Sweden.

^c Institute for the Psychology of Elite Performance, College of Health and Behavioural
Sciences, Bangor University, Bangor, Gwynedd, UK.

Corresponding Author:

Dr. Helen G. Hanstock

Swedish Winter Sports Research Centre

Department of Health Sciences

Mid Sweden University

831 40 Östersund

SWEDEN

Email: helen.hanstock@miun.se

Telephone: + 46 73 060 22 02

Word count (Abstract): 150/150

Word count (Body text): 1820/1500

26 **Key Words:**
27 Biomarkers
28 Tears
29 Noninvasive measures
30 Immunity
31 Stress reactivity
32 Immune reactivity
33 State anxiety
34 Acute stress
35 Psychological stress

Abstract

Tear secretory immunoglobulin-A (SIgA) is a putative biomarker of common-cold risk with potential utility in non-invasive diagnostics. As SIgA secretion at the ocular surface is under strong autonomic control, we investigated the relationship between HR reactivity and tear SIgA responses to novel experiential stress. Thirty-two healthy participants undertook a 60-second zip-line ride to evoke acute stress and a seated-rest control trial in a randomised-crossover design. We recorded heart rate (HR) continuously and collected unstimulated tear samples 5-min-pre-, 2-min-post- and 20-min-post-stress/control. Stress increased HR and state anxiety whereas tear SIgA concentration decreased 44% post-stress vs. control. Higher peak HR values during stress uniquely explained 21% of the variance in tear SIgA reactivity to stress ($p < .01$); high HR reactors displayed greater decreases in tear SIgA concentration. We conclude that physiological arousal increases immune reactivity to acute stress and highlight tear SIgA as a minimally-invasive, physiologically relevant biomarker of immune reactivity.

Introduction

Mucosal secretions are an attractive medium for the repeated, non-invasive assessment of endocrine, immune and inflammatory responses to stress (Papacosta & Nassis, 2011; Slavish, Graham-Engeland, Smyth, & Engeland, 2015). Secretory immunoglobulin-A (SIgA) provides a direct measure of immune competence due to its antimicrobial actions at the mucosal epithelia (Brandtzaeg, 2013). Low salivary SIgA levels have been highlighted as a risk factor for upper respiratory illness in athletes (Gleeson et al., 2012; Neville, Gleeson, & Folland, 2008) and the general population (Jemmott & McClelland, 1989; Volkmann & Weekes, 2006).

Several previous studies of mucosal immune responses to acute stressors have utilised salivary SIgA as a biomarker of immune reactivity to acute laboratory stressors (Benham, 2007; Bosch et al., 2001; Bosch, de Geus, Veerman, Hoogstraten, & Nieuw Amerongen, 2003; Campisi, Bravo, Cole, & Gobeil, 2012) and longer-term naturalistic stress (Engeland et al., 2016; Phillips et al., 2006; Volkmann & Weekes, 2006). However, the tear fluid offers an alternative, minimally-invasive medium to assess immune function. Transmission of upper respiratory tract infections (URTI) has been demonstrated at the ocular surface (Bischoff, Reid, Russell, & Peters, 2011) whereas oral transmission of URTI may be less common (Hendley & Gwaltney, 1988). It is likely that the tear fluid plays an important role in host defence and indeed recent evidence suggests that tear fluid SIgA can outperform salivary SIgA to assess URTI risk (Hanstock et al., 2016). Tear SIgA has been shown to decrease immediately after prolonged exercise (Hanstock et al., 2016), but the effect of acute stress on this putative immune biomarker remains unexplored.

Immune reactivity to acute experiential stress has been demonstrated in first-time skydivers (Schedlowski et al., 1993) and bungee jumpers (van Westerloo et al., 2011). These activities increase state anxiety (Hare, Wetherell, & Smith, 2013), activate sympathoadrenal-medullary and hypothalamic-pituitary-adrenal stress responses (Chatterton, Vogelsong, Lu, & Hudgens, 1997). Acute experiential stress may acutely activate cellular immune parameters, for example by mobilising NK cells (Schedlowski et al., 1993); a finding that has been mirrored in numerous studies employing acute laboratory-based stressors (Segerstrom & Miller, 2004), but may also inhibit innate immune function (van Westerloo et al., 2011). Individual differences in stress-induced sympathetic activation can predict the magnitude of cellular immune responses to acute laboratory stressors (Manuck, Cohen, Rabin, Muldoon, & Bachen, 1991; Marsland, Bachen, Cohen, Rabin, & Manuck, 2002). Given that secretion of SIgA at the ocular surface is under strong autonomic control (Dartt, 2009) it is likely that tear

SIgA reactivity to stress will correlate with other autonomic responses such as the heart rate (HR) response to stress. Thus, our aim was to investigate the relationship between HR, state anxiety and tear SIgA responses to a novel experiential stressor.

Method

Participants

Thirty-two healthy adults (17 males, 15 females) aged 23 years (SD = 4 years) provided informed consent to participate in the study. Participants had no previous experience of the stressor and avoided alcohol, caffeine, over-the-counter medication and heavy exercise for 24 h preceding experimental trials. No participants self-reported URTI symptoms during the 4 weeks prior to the study.

Experimental procedures

Participants completed two experimental trials on consecutive days in a randomised-crossover design. The stress trial involved a ride on a 1.6 km Zip-line (ZipWorld Velocity, Gwynedd, UK), lasting approximately 60 s. Participants wore a transparent plastic eye mask to prevent watering of the eyes during the ride. Trained instructors attached participants' safety harness to the line in a suspended prone position. Participant's movement was minimal in the suspended position and no physical effort was required to complete the task. During the control trial, participants sat quietly in the laboratory for 20 min. We recorded heart rate (HR) continuously in both trials (FT7, Polar Electro, Kempele, Finland) so that peak HR during stress (HR_{peak}) could be detected. Two participants' HR monitors recorded incomplete data and were excluded from HR-based analyses. To assess state anxiety, participants completed form Y1 of the State-Trait Anxiety Inventory (STAI-Y1; Spielberger, 1983) 5 min before each trial.

Sample collection, handling and analysis

We collected tear samples at 5-min-pre, 2-min-post and 20-min-post stress onset and at the same times of day during the control trial using methods previously described (Hanstock et al., 2016). Briefly, tear fluid collected from the inferior marginal tear strip via glass microcapillary pipette was transferred to a pre-weighed microcentrifuge tube and refrigerated. At 3 h post-collection, samples were weighed to 0.01 mg, diluted 1:99 in phosphate-buffered saline and frozen at -80°C. We demonstrated stability of SIgA-C in tear samples after 3 hours refrigeration in a pilot study (see Supplementary Material). After thawing, we used an enzyme-linked immunosorbent assay to determine tear SIgA-C in duplicate (Salimetrics, PA, USA; intra-assay CV = 1.6%). We calculated SIgA secretion rate (SIgA-SR) by multiplying tear flow rate (sample mass/collection time) by SIgA-C.

Statistical analyses

We performed statistical analyses using SPSS (v24, IBM, New York, USA) and GraphPad Prism (v5, San Diego, USA). With power 0.8 and alpha 0.05, we estimated a sample size of 32 participants for a model with three predictors to detect a large f^2 effect size of 0.4 (G*Power 3.1.9, Germany). Tear SIgA-C and SIgA-SR displayed log-normal distributions and were log-transformed before analysis. The efficacy of the zip-line ride to increase state anxiety and HR was assessed using paired t-tests; effect sizes are Cohen's d . Two-way repeated-measures ANOVA was used to explore the influence of stress on SIgA-C and SIgA-SR. Reactivity effects were explored using hierarchical linear regression. We defined tear SIgA reactivity as the difference in log-transformed values (\log_2 fold-change) between the control condition and 2-min-post-stress to give equal weighting to increases and decreases from control values in the regression analysis.

Results

Physiological and psychological responses to stress.

Peak HR during the zip-line ride was higher than mean HR during seated rest (Table 1); we defined this difference as Δ HR. Prior to the zip-line ride state anxiety increased compared to control (Table 1); we defined this difference as Δ STAI-Y1.

Table 1. Efficacy of zip-line protocol to increase HR and state anxiety.

	Stress Trial		Control Trial		Statistics			
	Mean Peak	SD	Mean	SD	<i>t</i>	<i>df</i>	<i>p</i>	<i>d</i>
Heart rate (bpm)	126	21	73	9	15.01	31	<.001	3.45
STAI-Y1 score	41	14	28	7	5.88	29	<.001	1.19

Effect of stress on tear SIgA-C and SIgA-SR.

Repeated-measures ANOVA revealed that tear SIgA-C decreased during the stress trial (time * trial interaction effect: $F(2,62) = 4.58, p = .01$; Fig 1a); Tukey's HSD revealed a reduction in SIgA-C at 2-min-post-stress compared to 5-min-pre-stress and lower SIgA-C during stress vs. control at all time points. At 2-min-post-stress, 28 of 32 participants' SIgA-C was lower than control, with a 44% mean decrease (SD = 36%, $d = 1.23$). There was a trend towards decreased SIgA-SR throughout the stress trial (main effect of trial: $F(1,31) = 3.37, p = .08$, Fig 1b).




Figure 1. Tear SIgA-C and SIgA-SR responses to stress and control. Mean \pm SD. Grey shade represents zip-line ride duration. Significant difference from 5-min-pre: *, $p < .05$, **, $p < .01$; ##, between trials, $p < .01$

Heart rate, state anxiety and tear SIgA reactivity to stress.

We used hierarchical linear regression to determine the relationship between stress reactivity and tear SIgA-C reactivity to stress. We entered participants' sex into the regression model first, followed by Δ HR at Step 2 and Δ STAI-Y1 at Step 3. Collinearity statistics were within accepted ranges. At Step 2 addition of Δ HR was able to significantly explain SIgA-C reactivity ($F(2,27) = 5.67, p = .009$), but addition of Δ STAI-Y1 at step 3 did not improve the model further (Table 2). No significant relationships were found between sex, Δ HR or Δ STAI-Y1 and SIgA-SR reactivity to stress ($F(3,26) = .77, p = .52$).

Table 2. Hierarchical linear regression reveals Δ HR as a significant explanatory variable for the tear SIgA-C response to stress. **, $p < .01$.

	Coefficients			Model		Change statistics		
	B	SE	β	R^2	F	df	ΔR^2	p
1				.090	2.78	1, 28	-	.106
(Constant)	-.294	.565	-					
Sex	-.622	.373	-.301					
2				.296	5.27	1, 27	.205	.009**
(Constant)	.655	.609	-					
Sex	-.341	.349	-.165					
Δ HR	-.025	.009	-.473**					
3				.317	0.28	1, 26	.022	.372
(Constant)	.683	.612	-					
Sex	-.330	.350	-.160					
Δ HR	-.030	.010	-.553**					
Δ STAI-Y1	.015	.016	.167					

Discussion

This study is the first to explore the effect of acute psychological stress on ocular immune parameters, and provides preliminary validation of tear SIgA-C as a biomarker of immune reactivity to acute stress. We observed that the zip-line protocol produced marked elevations of HR and state anxiety, and decreased tear SIgA-C throughout the duration of the stress trial. Participants with the greatest HR responses to the stressor tended to exhibit

greater decreases in tear SIgA post-stress. These observations support a role for physiological arousal in determining tear SIgA-C reactivity to stress.

During the stress trial, SIgA-C was lowest immediately post-stress, but was lower than control throughout, from 5-min before to 20-min after the zip line ride. That we did not blind participants to the stressor in advance likely caused anticipatory stress accounting for the lower tear SIgA-C at 5-min-pre; together with the lower tear SIgA-C at 20-min-post indicates that the salient influence of the stressor extends beyond 60 s duration of the zip line ride. The magnitude of the decrease in tear SIgA-C post-stress was a little smaller than previously reported decreases in tear SIgA-C following 2 h moderate-intensity exercise (-44% vs. -57%; Hanstock et al., 2016). These observations further support a role for physiological arousal, as occurs during exercise, in mediating the tear SIgA response to stress. Since the lacrimal gland secretions are primarily under parasympathetic control (Dartt, 2009), we speculate that the decrease in tear SIgA-C may arise as a result of the parasympathetic withdrawal that typically occurs during acute stress (Brindle, Ginty, Phillips, & Carroll, 2014). A limitation of this study was that we did not assess autonomic balance, but future studies could explore the relationship between autonomic activity and tear SIgA secretion in humans.

Tear SIgA-C has been previously highlighted as a potential biomarker of common cold risk (Hanstock et al., 2016). As the decrease in tear SIgA-C post-stress in the present study (-44%) was of greater magnitude than the 34% decrease in tear SIgA-C reported during the week before upper respiratory illness (Hanstock et al., 2016), the SIgA-C response to stress in the present study may have been of sufficient magnitude to compromise host defence in some of the higher reactors. These observations are consistent with the reactivity hypothesis which proposes that extremely high or low stress reactivity could exacerbate day-to-day fluctuations in immune function, increase susceptibility to opportunistic infections

(Cacioppo et al., 1998) and indicate poor states of long-term health (Lovallo, 2011). It has also been suggested that stress reactivity is a trainable trait and that lifestyle interventions such as exercise training (Forcier et al., 2006; Klaperski, von Dawans, Heinrichs, & Fuchs, 2014; von Haaren et al., 2016) and mindfulness meditation (Hoge et al., 2013) could attenuate stress reactivity, thus may have potential to improve health-related outcomes. Thus, future work is warranted to explore the influence of repeated daily hassles and subsequently lifestyle interventions on tear immunological responses to stress.

Here we demonstrate in a field-based study that tear SIgA-C is responsive to acute stress and that participants with higher HR reactivity display greater decreases in tear SIgA-C. This proof-of-concept study paves the way for future studies to examine tear SIgA responses to controlled laboratory stressors and naturalistic chronic stress. Characterising tear SIgA responses to acute and prolonged stress is warranted because the ocular surface is an important point of entry for pathogens that cause URTI (Bischoff et al., 2011) and because tear fluid is gaining interest as a medium from which to assess biomarkers (Farandos, Yetisen, Monteiro, Lowe, & Yun, 2015; Hagan, Martin, & Enríquez-de-Salamanca, 2016). If tear biomarkers are able to reliably predict health-related outcomes, wearable biosensors such as “smart” contact lenses could afford consumers the opportunity to self-monitor changes in immune status alongside other biomarkers of stress and health.

Acknowledgements

Funding: Helen Hanstock’s PhD was supported by a Bangor University 125th Anniversary Studentship.

We would like to thank the participants for their time and co-operation as well as Dr.

Matthew Fortes, Matthew Singleton, Daniel Kashi, Mark Ward, Xin Hui Aw Yong, Alex

Carswell, Karen Thomas, Larissa Gibson-Smith, Oliver Grounsell and Paul Gray for their assistance with data collection, and ZipWorld Velocity for allowing the study to take place.

References

- Benham, G. (2007). The shape of stress: the use of frequent sampling to measure temporal variation in S-IgA levels during acute stress. *Stress and Health*, 23(5), 295–301. <http://doi.org/10.1002/smi.1150>
- Bischoff, W. E., Reid, T., Russell, G. B., & Peters, T. R. (2011). Transocular entry of seasonal influenza-attenuated virus aerosols and the efficacy of N95 respirators, surgical masks, and eye protection in humans. *Journal of Infectious Diseases*, 204(2), 193–99. <http://doi.org/10.1093/infdis/jir238>
- Bosch, J. A., de Geus, E. J. C., Veerman, E. C. I., Hoogstraten, J., & Nieuw Amerongen, A. V. (2003). Innate secretory immunity in response to laboratory stressors that evoke distinct patterns of cardiac autonomic activity. *Psychosomatic Medicine*, 65(2), 245–258. <http://doi.org/10.1097/01.PSY.0000058376.50240.2D>
- Bosch, J. A., de Geus, E. J., Kelder, A., Veerman, E. C., Hoogstraten, J., & Amerongen, A. V. (2001). Differential effects of active versus passive coping on secretory immunity. *Psychophysiology*, 38(5), 836–846. <http://doi.org/10.1111/1469-8986.3850836>
- Brandtzaeg, P. (2013). Secretory IgA: Designed for anti-microbial defense. *Frontiers in Immunology*, 4(8), 1–17. <http://doi.org/10.3389/fimmu.2013.00222>
- Brindle, R. C., Ginty, A. T., Phillips, A. C., & Carroll, D. (2014). A tale of two mechanisms: A meta-analytic approach toward understanding the autonomic basis of cardiovascular reactivity to acute psychological stress. *Psychophysiology*, 51(10), 964–976. <http://doi.org/10.1111/psyp.12248>
- Cacioppo, J., Berntson, G., Malarkey, W., Kiecolt-Glaser, J., Sheridan, J., Poehlmann, K., ... Glaser, R. (1998). Autonomic, neuroendocrine, and immune responses to psychological stress: The reactivity hypothesis. *Annals of the New York Academy of Sciences*, 664–673. <http://doi.org/10.1111/j.1749-6632.1998.tb09605.x>
- Campisi, J., Bravo, Y., Cole, J., & Gobeil, K. (2012). Acute psychosocial stress differentially influences salivary endocrine and immune measures in undergraduate students. *Physiology and Behavior*, 107(3), 317–321. <http://doi.org/10.1016/j.physbeh.2012.09.003>
- Chatterton, R. T., Vogelsong, K. M., Lu, Y. C., & Hudgens, G. A. (1997). Hormonal responses to psychological stress in men preparing for skydiving. *Journal of Clinical Endocrinology and Metabolism*, 82(8), 2503–2509. <http://doi.org/10.1210/jcem.82.8.4133>
- Dartt, D. A. (2009). Neural regulation of lacrimal gland secretory processes: relevance in dry eye diseases. *Progress in Retinal and Eye Research*, 28(3), 155–77. <http://doi.org/10.1016/j.preteyeres.2009.04.003>
- Engeland, C. G., Hugo, F. N., Hilgert, J. B., Nascimento, G. G., Junges, R., Lim, H. J., ... Bosch, J. A. (2016). Psychological distress and salivary secretory immunity. *Brain, Behavior, and Immunity*, 52, 11–17. <http://doi.org/10.1016/j.bbi.2015.08.017>

- 271 Farandos, N. M., Yetisen, A. K., Monteiro, M. J., Lowe, C. R., & Yun, S. H. (2015). Contact
272 lens sensors in ocular diagnostics. *Advanced Healthcare Materials*, 4, 792–810.
273 <http://doi.org/10.1002/adhm.201400504>
- 274 Forcier, K., Stroud, L. R., Papandonatos, G. D., Hitsman, B., Reiches, M., Krishnamoorthy,
275 J., & Niaura, R. (2006). Links between physical fitness and cardiovascular reactivity and
276 recovery to psychological stressors: a meta-analysis. *Health Psychology*, 25(6), 723–
277 739. <http://doi.org/10.1037/0278-6133.25.6.723>
- 278 Gleeson, M., Bishop, N., Oliveira, M., McCauley, T., Tauler, P., & Muhamad, A. S. (2012).
279 Respiratory infection risk in athletes: association with antigen-stimulated IL-10
280 production and salivary IgA secretion. *Scandinavian Journal of Medicine & Science in
281 Sports*, 22(3), 410–7. <http://doi.org/10.1111/j.1600-0838.2010.01272.x>
- 282 Hagan, S., Martin, E., & Enríquez-de-Salamanca, A. (2016). Tear fluid biomarkers in ocular
283 and systemic disease: potential use for predictive, preventive and personalised medicine.
284 *The EPMA Journal*, 7(1), 15. <http://doi.org/10.1186/s13167-016-0065-3>
- 285 Hanstock, H. G., Walsh, N. P., Edwards, J. P., Fortes, M. B., Cosby, S. L., Nugent, A., ...
286 Yong, X. H. A. (2016). Tear fluid SIgA as a noninvasive biomarker of mucosal
287 immunity and common cold risk. *Medicine and Science in Sports and Exercise*, 48(3),
288 569–577. <http://doi.org/10.1249/MSS.0000000000000801>
- 289 Hare, O. A., Wetherell, M. A., & Smith, M. A. (2013). State anxiety and cortisol reactivity to
290 skydiving in novice versus experienced skydivers. *Physiology and Behavior*, 118, 40–
291 44. <http://doi.org/10.1016/j.physbeh.2013.05.011>
- 292 Hendley, J. O., & Gwaltney, J. M. (1988). Mechanisms of transmission of rhinovirus
293 infections. *Epidemiologic Reviews*, 10(2), 243–58.
294 <http://doi.org/10.1093/oxfordjournals.epirev.a036024>
- 295 Hoge, E. A., Bui, E., Marques, L., Metcalf, C. A., Morris, L. K., Robinaugh, D. J., ... Simon,
296 N. M. (2013). Randomized controlled trial of mindfulness meditation for generalized
297 anxiety disorder: effects on anxiety and stress reactivity. *The Journal of Clinical
298 Psychiatry*, 74(8), 786–792. <http://doi.org/10.4088/JCP.12m08083>
- 299 Jemmott, J. B., & McClelland, D. C. (1989). Secretory IgA as a measure of resistance to
300 infectious disease: Comments on Stone, Cox, Valdimarsdottir, and Neale. *Behavioral
301 Medicine*, 15(2), 63–71. <http://doi.org/10.1080/08964289.1989.9935153>
- 302 Klaperski, S., von Dawans, B., Heinrichs, M., & Fuchs, R. (2014). Effects of a 12-week
303 endurance training program on the physiological response to psychosocial stress in men:
304 a randomized controlled trial. *Journal of Behavioral Medicine*, 1118–1133.
305 <http://doi.org/10.1007/s10865-014-9562-9>
- 306 Lovallo, W. R. (2011). Do low levels of stress reactivity signal poor states of health?
307 *Biological Psychology*, 86(2), 121–128. <http://doi.org/10.1016/j.biopsycho.2010.01.006>
- 308 Manuck, S. B., Cohen, S., Rabin, B. S., Muldoon, M. F., & Bachen, E. A. (1991). Individual
309 differences in cellular immune response to stress. *Psychological Science*, 2(2), 111–115.
310 <http://doi.org/10.1111/j.1467-9280.1991.tb00110.x>
- 311 Marsland, A. L., Bachen, E. A., Cohen, S., Rabin, B., & Manuck, S. B. (2002). Stress,
312 immune reactivity and susceptibility to infectious disease. *Physiology & Behavior*,
313 77(4–5), 711–6. [http://doi.org/10.1016/S0031-9384\(02\)00923-X](http://doi.org/10.1016/S0031-9384(02)00923-X)
- 314 Neville, V., Gleeson, M., & Folland, J. P. (2008). Salivary IgA as a risk factor for upper
315 respiratory infections in elite professional athletes. *Medicine and Science in Sports and*

- Exercise*, 40(7), 1228–36. <http://doi.org/10.1249/MSS.0b013e31816be9c3>
- Papacosta, E., & Nassis, G. P. (2011). Saliva as a tool for monitoring steroid, peptide and immune markers in sport and exercise science. *Journal of Science and Medicine in Sport*, 14(5), 424–34. <http://doi.org/10.1016/j.jsams.2011.03.004>
- Phillips, A. C., Carroll, D., Evans, P., Bosch, J. A., Clow, A., Hucklebridge, F., & Der, G. (2006). Stressful life events are associated with low secretion rates of immunoglobulin A in saliva in the middle aged and elderly. *Brain, Behavior, and Immunity*, 20(2), 191–7. <http://doi.org/10.1016/j.bbi.2005.06.006>
- Schedlowski, M., Jacobs, R., Stratmann, G., Richter, S., Hädicke, A., Tewes, U., ... Schmidt, R. E. (1993). Changes of natural killer cells during acute psychological stress. *Journal of Clinical Immunology*, 13(2), 119–26. <http://doi.org/10.1007/BF00919268>
- Seegerstrom, S. C., & Miller, G. E. (2004). Psychological stress and the human immune system: a meta-analytic study of 30 years of inquiry. *Psychological Bulletin*, 130(4), 601–30. <http://doi.org/10.1037/0033-2909.130.4.60>
- Slavish, D. C., Graham-Engeland, J. E., Smyth, J. M., & Engeland, C. G. (2015). Salivary markers of inflammation in response to acute stress. *Brain, Behavior, and Immunity*, 44, 253–269. <http://doi.org/10.1016/j.bbi.2014.08.008>
- Spielberger, C. (1983). *Manual for the State-Trait Anxiety Inventory*. Palo Alto, CA: Mind Garden.
- van Westerloo, D. J., Choi, G., Löwenberg, E. C., Truijen, J., de Vos, A. F., Endert, E., ... van der Poll, T. (2011). Acute stress elicited by bungee jumping suppresses human innate immunity. *Molecular Medicine*, 17(3–4), 180–188. <http://doi.org/10.2119/molmed.2010.00204>
- Volkman, E. R., & Weekes, N. Y. (2006). Basal SIgA and cortisol levels predict stress-related health outcomes. *Stress and Health*, 22(1), 11–23. <http://doi.org/10.1002/smi.1077>
- von Haaren, B., Ottenbacher, J., Muenz, J., Neumann, R., Boes, K., & Ebner-Priemer, U. (2016). Does a 20-week aerobic exercise training programme increase our capabilities to buffer real-life stressors? A randomized, controlled trial using ambulatory assessment. *European Journal of Applied Physiology*, 116(2), 383–394. <http://doi.org/10.1007/s00421-015-3284-8>